

O I Line Emission in the Quasar PG 1116+215

Yoshiki MATSUOKA,^{1*} Shinki OYABU,^{2*} Yumihiko TSUZUKI,³ Kimiaki KAWARA,¹ and Yuzuru YOSHII¹

¹*Institute of Astronomy, School of Science, The University of Tokyo*

2-21-1, Osawa, Mitaka, Tokyo 181-0015

matsuoka@ioa.s.u-tokyo.ac.jp

²*Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency*

3-1-1, Yoshinodai, Sagami-hara, Kanagawa 229-8510

³*Institute for Cosmic Ray Research, The University of Tokyo 5-1-5, Kashiwanoha, Kashiwa, Chiba 277-8582*

(Received 2005 April 28; accepted 2005 May 20)

Abstract

By observing the near-infrared spectrum of the quasar PG 1116+215 at $z = 0.176$ and combining with the HST/FOS spectrum, we obtained the relative strengths of three permitted O I lines ($\lambda 1304$, $\lambda 8446$, and $\lambda 11287$) in a quasar for the first time. The photon flux ratios of the O I lines of the quasar were compared with those previously measured in a Seyfert 1 and six narrow-line Seyfert 1s. No significant differences were found in the O I line flux ratios between the quasar and the other Seyferts, suggesting that the gas density in the O I and Fe II line-emitting regions in the quasar is of the same order as those in low-luminosity AGNs. It was also found that the line width of O I $\lambda 11287$ is significantly narrower than that of Ly α , which is consistent with O I and Fe II emission occurring in the partly ionized regions at the outermost portion of the broad-line region where velocities are small.

Key words: galaxies: active — galaxies: individual (PG 1116+215) — galaxies: quasars: emission lines — line: formation

1. Introduction

The permitted O I lines at 1304, 8446, and 11287 Å are common features in AGN (active galactic nuclei) and have been used to study the physical properties of BLR (broad-line region) clouds (e.g., Grandi 1980; Kwan & Krolik 1981; Rudy et al. 1989; Laor et al. 1997). As shown in the partial Grotrian diagram of O I atom in figure 1, the transition from the ground state $2p^3P$ to the excited state $3d^3D^0$ has an excitation energy whose wavelength is $\lambda = 1025.77$ Å, falling within the Doppler core of Ly β (1025.72 Å) for gas at 10^4 K. The excited state $3d^3D^0$ decays either back to the ground state, or by the emission of a $\lambda 11287$ photon, to the intermediate excited state $3p^3P$. The latter decays by a $\lambda 8446$ photon to the lower excited state $3s^3S^0$, which finally decays to the ground state by a $\lambda 1304$ photon. If Ly β pumping is the dominant process of O I line formation, the photon flux ratio of these three lines should be 1:1:1, and the photon flux ratio, especially between O I $\lambda 1304$ and $\lambda 8446$, can be used as a reddening indicator (Rudy et al. 1989; Laor et al. 1997; Rodríguez-Ardila et al. 2002b). It can also be expected that if the O I line emission is dominated by Ly β pumping, the O I line intensities should be used to measure the microturbulence parameter in BLR clouds, because Ly β pumping would become more efficient in a larger microturbulent velocity

field.

Several mechanisms that could modify the 1:1:1 photon flux ratio have been suggested, such as the collisional excitation of O I $\lambda 8446$ and the Balmer continuum absorption of O I $\lambda 1304$ (Grandi 1983). Rodríguez-Ardila et al. (2002b) found that in six of their seven low-luminosity AGNs, Ly β pumping is not the only mechanism responsible for the O I line emission, and the 1:1:1 photon flux ratio is altered. Their sample consisted of six NLS1s (narrow line Seyfert 1s) and one Seyfert 1. Before Rodríguez-Ardila et al. (2002b), I Zw 1, classified as the NLS1, is the only object where the three O I lines have been studied (Rudy et al. 1989, 2000; Laor et al. 1997). It is thus not clear whether the properties of the O I line emission found in low-luminosity AGNs can be applied to quasars. The O I line emission in quasars would differ from those of NLS1s, because it is generally considered that the gas density in BLR clouds is higher in NLS1s than in quasars, and thus the collisional processes are more dominant in NLS1s (Baldwin et al. 1988, 1996; Laor et al. 1997; Wilkes et al. 1999; Kuraszkiewicz et al. 2000). Investigating the O I line formation mechanisms would also provide knowledge about the physical properties of the Fe II emitting region, since the Fe II and O I lines are considered to be emitted in the same portion of BLR clouds (Rodríguez-Ardila et al. 2002a). Since Fe II emission is the strongest coolant in BLR clouds, it would give an important clue to better understand the radiative mechanisms in BLR gas.

In order to study the O I line formation mechanism in quasars, we have started a program of taking NIR(near-infrared) spectra of quasars. Combining the NIR spectra

* Visiting Astronomer, Kitt Peak National Observatory, National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation, USA.

with the HST/FOS ultraviolet spectra, we intend to analyze the three O I lines, i.e., O I λ 1304, O I λ 8446, and O I λ 11287. We present the ratios of the three O I lines of the quasar PG 1116+215 at $z=0.176$. This is the first quasar in which the three O I lines have been analyzed.

2. Observation and Reduction

2.1. Near-Infrared Spectroscopy

The NIR spectrum of PG 1116+215, covering $0.9 - 1.8\mu\text{m}$, including the redshifted O I λ 8446 and O I λ 11287 lines, was obtained using the longslit of FLAMINGOS (Florida Multi-object Imaging Near-IR Grism Observational Spectrometer; Elston 1998) on the KPNO (Kitt Peak National Observatory) 2.1m telescope on 2005 February 28. The detector is the Hawaii II 2048×2048 HgCdTe science-grade array, divided into four quadrants with 8 amplifiers each. A 4-pixel slit with a scale of $0''.606 \text{ pixel}^{-1}$ was placed on the object in the north-south direction under a seeing of $1''$. The first order of the *JH* grism was used, which gave a spectral resolution of 430 km s^{-1} . The object was shifted along the slit by $20''$ between exposures. The total integration time was 2400s, which consisted of eight 300s exposures. *J*-band photometry of the object was also performed for photometric calibration.

Data reduction was performed using IRAF (Image Reduction and Analysis Facility).¹ OH airglow lines were used to calibrate the wavelength scale. The flux scaling was twofold: (1) the A-type star SAO 81808 was observed to calibrate the relative response within the *J*-band, and (2) the standard star AS 20-0, having $J = 9.55 \pm 0.01$ mag (Hunt et al. 1998), was used to determine the *J*-band magnitude. A 9500K blackbody was fitted to the spectrum of SAO 81808 to obtain the sensitivity curve within the *J*-band. Both standard stars were observed at an airmass of 1.1, which is similar to that of the object. The total uncertainty in flux calibration is 5%. The final spectrum was re-binned into a $10 \text{ \AA}$ step, corresponding to a spectral resolution of 300 km s^{-1} .

2.2. Ultraviolet Spectroscopy

The HST (Hubble Space Telescope) ultraviolet spectrum of PG 1116+215 was observed on 1993 February 19 and 20 by using the FOS (Faint Object Spectrograph) with three gratings (G130H, G190H, and G270H), which cover $1200 - 3200 \text{ \AA}$ in the observed frame. The integration times were 8724s for G130H, 1878s for G190H, and 751s for G270H.

The HST spectrum used in this work was obtained from Evans & Koratkar (2004). They recalibrated the raw archival spectra using the latest algorithms and calibration data. Spectral data contaminated by intermittent noisy diodes and cosmic-ray events were identified manually and eliminated. They combined multiple observa-

tions of the same source, if available, in such a way that the resultant spectrum would have the highest possible signal-to-noise ratio. We removed prominent geocoronal and galactic absorption lines from the spectrum of PG 1116+215 by interpolating from both sides of the feature. The spectrum of PG 1116+215 was re-binned into a $1 \text{ \AA}$ step, corresponding to a spectral resolution of 300 km s^{-1} . The total uncertainty in the flux calibration was less than 5%.

2.3. Galactic and Internal Extinction

In order to correct the galactic reddening, we adopted $E_{B-V} = 0.095 \pm 0.015$ mag from the extinction map of the Milky Way based on the far-infrared emission observed by IRAS and COBE/DIRBE (Schlegel, Finkbeiner, & Davis 1998). The resolution of their extinction map is $6'.1$. Dereddening of our spectrum was performed by using a galactic extinction curve presented by Pei (1992).

We also sought the effect of the intrinsic extinction of PG 1116+215. Popović (2003) analyzed the HST/FOS spectrum of PG 1116+215, and found a flux ratio of $H\alpha/H\beta = 2.94 \pm 0.74$. When corrected for galactic reddening, as described above, the ratio became 2.70. Dong et al. (2005) showed that the flux ratio of the broad components of $H\alpha$ to $H\beta$ is $H\alpha/H\beta = 2.97 \pm 0.36$ for their 94 blue AGN samples. These AGN samples are considered to be free of intrinsic extinction. Judging from $H\alpha/H\beta$ of PG 1116+215, which is similar to those of these blue AGNs, the intrinsic extinction in PG 1116+215 appears to be little.

2.4. Variability

It is widely known that Seyferts and quasars are highly variable. Since the observation dates of the UV and NIR spectra are separated by more than 12 yrs, a possible variability effect should be carefully examined before comparing the O I lines in the UV and NIR wavelengths. $J = 13.59 \pm 0.03$ mag is found in the 2MASS (Two Micron All Sky Survey) database,² while $J = 12.96 \pm 0.03$ mag was obtained for our observation. It is thus clear that PG 1116+215 varied its brightness by about 0.6 mag since the 2MASS observation on 1998 February 2. However, because the O I lines are formed in the outermost portion of the BLR, it is expected that their variability is sufficiently small to be ignored, even if the power-law continuum, which is supposed to be direct light from the central source, shows significant variability. In fact, a monitoring observation of the well-known variable Seyfert NGC 5548 revealed that when the continuum varied its brightness by about three times, the Mg II line, which is emitted in the same region as the O I lines, varied by less than 6% (Dietrich & Kollatschny 1995). These arguments suggest that the variability has little effect on the O I line fluxes

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation, USA.

² This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation, USA.




Fig. 1. Partial Grotrian diagram of O I. The solid lines are used for permitted transitions, while the dashed lines are for semi-forbidden transitions.

of our data.

3. Results and Analysis

Figure 2 shows the UV spectrum of PG 1116+215 on the top panel and the NIR spectrum on the bottom. The three O I lines are clearly detected. Note that the signal-to-noise ratio of the NIR spectrum at the wavelength range shorter than $1\mu\text{m}$ is low because of the aberration at the edge of the detector. The O I $\lambda 11287$ line flux, which is free from blending with other lines, was measured by integrating all counts above the estimated continuum level in the wavelength range between $1.319\mu\text{m}$ and $1.338\mu\text{m}$ in the observed frame. The line width was measured by fitting a single Gaussian to the observed line. The FWHM of O I $\lambda 11287$ is $2000 \pm 200 \text{ km s}^{-1}$, and significantly narrower than that of Ly α ($\sim 5500 \text{ km s}^{-1}$). Rodríguez-Ardila et al. (2002a) showed that the line widths and profiles of Fe II, O I, and Ca II in their NLS1 samples are very similar, while they are significantly narrower than that of Pa β . Their results are consistent with the Fe II, O I, and Ca II lines occurring in the partly ionized regions,³ which are farther from the central ionizing source and have smaller velocities. Our narrow O I $\lambda 11287$ line width relative to Ly α implies that this geometrical configuration found in the NLS1s also applies to quasars.

Although O I $\lambda 8446$ is blended with the Ca II lines in the red wing, the blueward part of the line is almost free from its contamination. We thus fitted a single Gaussian to the line in the blueward part where the contamination by Ca II is not severe, and then measured the flux within




Fig. 2. UV spectrum taken with the HST/FOS shown on the top panel and the NIR spectrum on the bottom panel in the observed frame. The positions of the three O I lines, Ca II lines, and Si II $\lambda 1264$ line are indicated by vertical lines.

the fitted Gaussian.

The broad feature at $1304 \text{ \AA}$ results from the blending of an O I triplet at $\lambda 1302.17$, $\lambda 1304.86$, and $\lambda 1306.0$ and a Si II doublet at Si II $\lambda 1304$ and $\lambda 1309$. The total blend flux of the $\lambda 1304$ feature was measured by integrating all counts above the estimated continuum level between $1524 \text{ \AA}$ and $1548 \text{ \AA}$ in the observed frame. The O I and Si II lines are severely blended and de-blending with high spectral resolution is hopeless for quasars. However, the Si II doublet blended in the $1304 \text{ \AA}$ feature is expected to be accompanied by considerable Si II line emission at $1264 \text{ \AA}$ (Constantin et al. 2002). Actually, Si II $\lambda 1264$ is clearly detected in UV spectrum of PG 1116+215 (figure 2) and its flux is $44 \pm 5\%$ of the blend flux of the $1304 \text{ \AA}$ feature. The photoionization models by Kwan & Krolik (1979) and Netzer (1980) predict that the Si II doublet at $1304 \text{ \AA}$ and $1309 \text{ \AA}$ has $16 - 28\%$ of the flux of Si II $\lambda 1264$ (Dumont & Mathez 1981), implying that $7 - 12\%$ of the blend flux comes from the Si II doublet and $88 - 93\%$ from the O I triplet. Laor et al. (1997) found that $50 - 56\%$ of the blend flux is due to the O I triplet in I Zw 1 and Rodríguez-Ardila et al. (2002b) found that the average portion of O I flux contribution to the blend flux is 75% in their three NLS1s. In a sample of high-redshift quasars, Constantin et al. (2002) concluded that the $\lambda 1304$ feature is due only to O I lines, because Si II lines, which are

³ Note that O I, Fe II, and Ca II emitting regions should be heavily overlapped because of the similar ionization potentials, i.e., 13.6eV for O I, 16.2eV for Fe II, and 11.9eV for Ca II.

Table 1. Measured flux of O I $\lambda 1304$, $\lambda 8446$, and $\lambda 11287$ for PG 1116+215.

Line	Flux ($10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$)
O I $\lambda 1304$	27.4 ± 2.1
O I $\lambda 8446$	8.1 ± 1.7
O I $\lambda 11287$	4.6 ± 0.6

expected to be accompanied with Si III $\lambda 1304$ and $\lambda 1309$, are not seen. From the discussions presented above, we assumed that 90% of the $\lambda 1304$ of PG 1116+215 is due to O I lines. We emphasize that assuming the portion of O I lines as being 75% (Rodríguez-Ardila et al. 2002b) or 100% (Constantin et al. 2002) does not change our conclusions.

We list the measured flux of O I $\lambda 1304$, $\lambda 8446$, and $\lambda 11287$ in Table 1.

4. Mechanism of O I Line Formation

The photon flux ratios for PG 1116+215 of O I $\lambda 11287$ /O I $\lambda 8446$ (hereafter ROI_{ir}) and O I $\lambda 1304$ /O I $\lambda 8446$ (hereafter ROI_{uv}) are given in table 2 along with those for a Seyfert 1 and six NLS1s. The O I $\lambda 1304$ flux of I Zw 1 was obtained from Laor et al. (1997), while $\lambda 8446$ and $\lambda 11287$ were obtained from Rudy et al. (2000). ROI_{ir} and ROI_{uv} of other 5 NLS1s and a Seyfert 1 were obtained from Rodríguez-Ardila et al. (2002b). Note that while the quoted errors for PG 1116+215 and I Zw 1 are 1σ significant, those for other samples are 2σ significant.

The average value of ROI_{ir} for the NLS1s is 0.73 ± 0.22 , where the 1σ value, 0.22, reflects only the scatter from object to object, not including the errors measured for each object. ROI_{ir} of PG 1116+215 falls within this 1σ scatter of the NLS1 average. Even if we neglect Ton S180, which has a significantly larger value of ROI_{ir} than the other five NLS1s, the average value of ROI_{ir} for the NLS1s is 0.66 ± 0.15 and that of PG 1116+215 falls within the 1σ scatter. Thus, ROI_{ir} of PG 1116+215 is not significantly different from those of other NLS1 samples. In the same way, we can see no significant difference between the ROI_{ir} of the Seyfert 1 galaxy NGC 863 and those of the NLS1s.

ROI_{ir} of the AGN was first measured for I Zw 1 by Rudy et al. (1989). They found that ROI_{ir} of this object is equal to unity, and suggested that O I lines are formed by Ly β pumping. However, Rodríguez-Ardila et al. (2002b) found that ROI_{ir} of I Zw 1 falls below unity, using the spectrum with higher resolution and wider spectral coverage published by Rudy et al. (2000). They also examined the spectra of six NLS1s including I Zw 1, and one Seyfert 1, to find that in six of their seven samples, ROI_{ir} are significantly below unity. Our result for PG 1116+215 indicated that this trend is also present in quasars. They discussed that the collisional excitation to the upper level of the O I $\lambda 8446$ transition enhances the strength of the O I $\lambda 8446$ relative to that of the O I $\lambda 11287$, so that ROI_{ir} falls below unity. This conclusion was drawn from the presence of O I $\lambda 7774$ in 1H 1934–063, whose relative strength to

O I $\lambda 8446$ is consistent with the combination of Ly β pumping and collisional excitation mechanisms.

Rodríguez-Ardila et al. (2002b) ruled out continuum fluorescence as the additional mechanism from the lack or weakness of other O I lines, such as $\lambda 7002$, $\lambda 7254$, and $\lambda 13165$, which should be present if continuum fluorescence is at work (Grandi 1980). Measuring the strengths of O I $\lambda 6048$, $\lambda 7774$, and $\lambda 7990$, which are very weak relative to that of $\lambda 8446$, they also suggested that the contribution to O I $\lambda 8446$ by recombination is no larger than a few percents.

The average value of ROI_{uv} for the NLS1s is 0.33 ± 0.30 , adopting the upper limit value for 1H 1934–063. ROI_{uv} of PG 1116+215 and NGC 863 fall within this 1σ scatter of the NLS1s average, which means that the ROI_{uv} values of these two objects are not significantly different from those of other NLS1 samples. These values are significantly below unity. Kwan & Krolik (1981) showed that the intrinsic ROI_{uv} is 0.76, rather than 1, due to the collisional excitation of O I $\lambda 8446$ and the Balmer continuum absorption of O I $\lambda 1304$. Grandi (1983) found that O I $\lambda 1304$ could also be destroyed by collisional de-excitation of the O I $\lambda 1304$ transition and de-excitation of the upper term of O I $\lambda 1304$ via O I] $\lambda 1641$ and $\lambda 2324$ line emission. He calculated the fraction of populations of $3s \ ^3S^0$ that result in the observable $\lambda 1304$, and found that up to half of the $\lambda 1304$ photon could be converted to O I] $\lambda 1641$ before it leaves the emission-line cloud. However, Laor et al. (1997) found that no strong line is detected at 1641 Å for I Zw 1, and that such a line cannot add more than $\sim 30\%$ to the observed $\lambda 1304$ flux. Such an investigation was possible for I Zw 1 because the He II $\lambda 1640$ line, which is usually the dominant feature around 1641 Å, is blueshifted by about 10 Å in the I Zw 1 spectrum. Thus, they ruled out this mechanism as the main contributor to destroy the O I $\lambda 1304$ photon. Consequently, the major mechanism of O I $\lambda 1304$ destruction is the Balmer continuum absorption or collisional de-excitation, or both.

5. Discussion

Laor et al. (1997) showed that the relative strength of C III] $\lambda 1909$ of I Zw 1 is significantly lower than that observed in the typical AGN. They argued that this suppression of C III] $\lambda 1909$ may imply that the typical BLR density in I Zw 1 is about an order of magnitude larger than in the typical AGN. This suppression of C III] $\lambda 1909$ is also seen in other narrow-line quasars (Baldwin et al. 1988, 1996; Wilkes et al. 1999; Kuraszkiewicz et al. 2000), suggesting that the high density of the BLR gas is common in this type of AGN. However, it is not clear whether these arguments are applicable to the region where Fe II and O I lines are emitted. Kuraszkiewicz et al. (2000) showed that while line flux ratios of Ly α , C IV $\lambda 1549$, Si IV $\lambda 1440$, C III] $\lambda 1909$, and Si III] $\lambda 1892$ observed in the NLS1s can be explained by 10-times lower ionization parameter and a few-times (<10) higher densities than the normal AGN, the Mg II line strength cannot be explained by these parameter values. Since the Fe II and O I lines are considered

Table 2. Measured ROI_{uv} and ROI_{ir} for PG 1116+215, one Seyfert1, and six NLS1s.

Object	Type	M_B^*	$\text{ROI}_{\text{uv}}^{\dagger,\ddagger}$	$\text{ROI}_{\text{ir}}^{\dagger,\ddagger}$	References ^{II}
PG 1116+215	quasar	-24.7	0.52 ± 0.12	0.76 ± 0.19	—
NGC 863	Seyfert1	-20.9	0.26 ± 0.04	0.55 ± 0.08	1
1H 1934-063	NLS1	-17.1	< 0.19	0.64 ± 0.05	1
Ark 564	NLS1	-20.3	0.07 ± 0.01	0.82 ± 0.03	1
Mrk 335	NLS1	-21.0	0.20 ± 0.02	0.64 ± 0.05	1
Mrk 1044	NLS1	-19.5	0.43 ± 0.07	0.42 ± 0.05	1
Ton S180	NLS1	-22.6	0.91 ± 0.15	1.08 ± 0.16	1
I Zw 1*	NLS1	-22.7	0.19 ± 0.02	0.76 ± 0.11	2, 3

* These values were obtained by modifying the entries in the catalog presented by Véron-Cetty & Véron (2003), assuming cosmological constants as $H_0=70\text{km/s}$, $\Omega_M=0.3$, and $\Omega_\Lambda=0.7$. Originally assumed values are $H_0=50\text{km/s}$ and $q_0=0$.

[†] ROI_{uv} and ROI_{ir} are defined as the photon flux ratio of $\lambda 1304/8446$ and $\lambda 11287/8446$.

[‡] The quoted errors are 1σ significant for PG 1116+215 and I Zw 1, while those of other samples are 2σ significant.

* We assumed that 53% of the $\lambda 1304$ feature is due to O I lines, which is the average portion of optically thick and thin cases presented by Laor et al. (1997).

^{II} References.— (1) Rodríguez-Ardila et al. (2002b). (2) Laor et al. (1997). (3) Rudy et al. (2000).

to be formed in the same region as the Mg II line (see, e.g., Kwan & Krolik 1981), the physical properties in the Fe II emitting region could be quite different from the C III] emitting region.

Three O I lines presented here ($\lambda 1304$, $\lambda 8446$, and $\lambda 11287$) provide a good indicator of the gas density in the Fe II emitting region. This is because these O I line emissions are seriously affected by several mechanisms that are sensitive to the gas density, as described above. Our observation of these three O I lines in a quasar enables, for the first time, to investigate whether there is a significant difference in the gas density of Fe II emitting cloud between NLS1s and quasars. The results are quite different from the situation indicated by the C III] $\lambda 1909$ line. As described above, there are no significant differences in ROI_{ir} and ROI_{uv} among the NLS1s, the Seyfert 1, and the quasar. This indicates that the physical properties of O I emitting cloud affect the O I line formation in the similar way in these three types of AGN. In other words, the collisional processes are working to a similar extent in the NLS1s, the Seyfert1, and the quasar. Thus, our O I observations did not find any significant differences among these types of AGN in the gas density in the outermost portion of the BLR where the Fe II and O I lines are emitted. This result would provide some clues for modeling the environment of the Fe II emitting cloud in NLS1s, Seyfert 1s, and quasars.

NLS1s are known as strong Fe II emitters. The efficiency of Fe II emission is sensitive to several physical parameters other than the gas density, in which the most sensitive parameters are the ionization parameter, microturbulence, and input spectral energy distribution (Netzer & Wills 1983; Wills, Netzer, & Wills 1985; Verner et al. 1999, 2003). Our results suggest that the Fe II emission enhancement in the NLS1s may not be caused by the high density of the BLR gas. If this is the case, some of the other parameters should be quite different between NLS1s

and quasars.

6. Summary

We performed NIR spectroscopy of the quasar PG 1116+215. By combining the NIR spectrum with the UV spectrum taken with the HST/FOS, we obtained three O I lines ($\lambda 1304$, $\lambda 8446$, and $\lambda 11287$). We found that the line width of O I $\lambda 11287$ is narrower than that of Ly α , which is consistent with O I and Fe II emission occurring in the partly ionized regions at the outermost portion of the BLR. We also found that the photon flux ratio of the three O I lines significantly deviate from 1:1:1, the expected ratio in the case of pure Ly β pumping formation. This strongly suggests the contribution of mechanisms other than Ly β pumping to the O I line formation/destruction, for which the best candidates are the density-sensitive processes. Furthermore, the obtained photon flux ratio for PG 1116+215 is not significantly different from those of the NLS1s and the Seyfert 1. This indicates that the gas density in Fe II and O I emitting regions are not significantly different among NLS1s, Seyfert 1s, and quasars, although a larger gas density in the NLS1s is indicated by the C III] $\lambda 1909$ strengths. We also suggest that the physical parameters other than the gas density should be quite different between NLS1s and quasars to account for NLS1s as strong Fe II emitters.

We are grateful to the referee, Jack Baldwin, for useful comments to improve this manuscript. We thank the staff of KPNO for technical support and assistance with the observation. We would also like to thank H. Fukushi for her help. The trip of YM and SO to Tucson was financially supported by Research Center for the Early Universe, The University of Tokyo.

References

- Baldwin, J. A., Ferland, G. J., Korista, K. T., Carswell, R. F., & Hamann, F. 1996, *ApJ*, 461, 664
- Baldwin, J. A., McMahon, R., Hazard, C., & Williams, R. E. 1988, *ApJ*, 327, 103
- Constantin, A., Shields, J. C., Hamann, F., Foltz, C. B., & Chaffee, F. H. 2002, *ApJ*, 565, 50
- Dietrich, M., & Kollatschny, W. 1995, *A&A*, 303, 405
- Dong, X.-B., Zhou, H.-Y., Wang, T.-G., Wang, J.-X., Li, C., & Zhou, Y.-Y. 2005, *ApJ*, 620, 629
- Dumont, A. M., & Mathez, G. 1981, *A&A*, 102, 1
- Elston, R. 1998, *Proc. SPIE*, 3354, 404
- Evans, I. N., & Koratkar, A. P. 2004, *ApJS*, 150, 73
- Grandi, S. A. 1980, *ApJ*, 238, 10
- Grandi, S. A. 1983, *ApJ*, 268, 591
- Hunt, L. K., Mannucci, F., Testi, L., Migliorini, S., Stanga, R. M., Baffa, C., Lisi, F., & Vanzi, L. 1998, *AJ*, 115, 2594
- Kuraszkiewicz, J., Wilkes, B. J., Czerny, B., & Mathur, S. 2000, *ApJ*, 542, 692
- Kwan, J., & Krolik, J. H. 1979, *ApJ*, 233, L91
- Kwan, J., & Krolik, J. H. 1981, *ApJ*, 250, 478
- Laor, A., Jannuzi, B. T., Green, R. F., & Boroson, T. A. 1997, *ApJ*, 489, 656
- Netzer, H. 1980, *ApJ*, 236, 406
- Netzer, H., & Wills, B. J. 1983, *ApJ*, 275, 445
- Pei, Y. C. 1992, *ApJ*, 395, 130
- Popović, L. C. 2003, *ApJ*, 599, 140
- Rodríguez-Ardila, A., Viegas, S. M., Pastoriza, M. G., & Prato, L. 2002a, *ApJ*, 565, 140
- Rodríguez-Ardila, A., Viegas, S. M., Pastoriza, M. G., Prato, L., & Donzelli, C. J. 2002b, *ApJ*, 572, 94
- Rudy, R. J., Mazuk, S., Puetter, R. C., & Hamann, F. 2000, *ApJ*, 539, 166
- Rudy, R. J., Rossano, G. S., & Puetter, R. C. 1989, *ApJ*, 342, 235
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Verner, E. M., Bruhweiler, F., Verner, D., Johansson, S., & Gull, T. 2003, *ApJ*, 592, L59
- Verner, E. M., Verner, D. A., Korista, K. T., Ferguson, J. W., Hamann, F., & Ferland, G. J. 1999, *ApJS*, 120, 101
- Véron-Cetty, M.-P., & Véron, P. 2003, *A&A*, 412, 399
- Wilkes, B. J., Kuraszkiewicz, J., Green, P. J., Mathur, S., & McDowell, J. C. 1999, *ApJ*, 513, 76
- Wills, B. J., Netzer, H., & Wills, D. 1985, *ApJ*, 288, 94